Space Weather Technical Forecast

Issued on Friday, 17 May 2024 at 13:12 Local

This technical forecast provides a four day assessment of space weather events. The probabilities stated below are for reaching or exceeding the given levels. For more information about space weather impacts please see the Met Office Space Weather Scales https://www.metoffice.gov.uk/weather/learn-about/space-weather/uk-scales

Space Weather Forecast Headline: Likely chance of isolated M-class flares, and Slight chance of isolated X-class flares throughout. Decreasing Chance of renewed S1/Minor Solar Radiation storming Day 1 (17 May).

Analysis of Space Weather Activity over past 24 hours

Solar Activity: Solar activity has been Low in the past 24 hours, with a peak C8.7 flare observed at 17/0146 UTC from AR3679. There are currently 14 sunspots on the visible disc, 13 of which are numbered. Any further activity from Ex-AR3664 is likely to be completely obscured as it has rotated onto the far-side. The largest region on the disc is AR3685 (Ehi/Beta), which lies close to the southeast limb. This region is composed of a number of large spots which may be a closely connected collection of different regions, magnetic looping is visible across the region and as such it remains classed as one large sunspot group. A new region continues to emerge from the southeast limb, just to the north of AR3685. At present it looks to be a single moderately large spot and is initially classed as an Hax/Alpha type. AR3679 (Esi/Beta-Gamma) is the second largest region, currently approaching the south central disc. It has also has been the source of most of the recent flare activity, with some further trailer spot emergence evident. AR3683 (Dsi/Beta-Gamma) in the southwest is of modest magnetic complexity and continues to show spot development/emergence. The other sunspot regions are generally smaller, simpler and stable, or in slight decay.

No Earth-directed CMEs were identified in available imagery.

Solar Wind / Geomagnetic Activity: Any residual CME effects at the start of the period, soon waned. Solar winds declined then settled in the range 370-420km/s. Density of the solar wind was initially above average, near 27 ppcc, but quickly fell back to average levels. Interplanetary Magnetic Field (IMF), Bt, was occasionally moderate, peak 10nT. The north-south component Bz varied between+10nT to -7nT, but was largely positive (northward). Phi angle was predominantly positive (away from the Sun) with brief periods of negative (towards the Sun) orientation, although there was an extended period of negative 16/1500-2100 UTC. Geomagnetic activity was initially Unsettled to Active (Kp 3 to 4), but fell to Quiet levels (Kp 1-2) from 16/1800 UTC.

Energetic Particles / Solar Radiation: High energy proton flux (greater than 10 MeV) by GOES-18 was initially at S1/Minor Radiation Storm levels, but declined to slightly elevated levels from around 16/1450 UTC and has since seen a continued slow decline. High energy electron flux (greater than 2MeV) observed by GOES16 was at predominantly background levels. The associated 24-hour fluence has remained below the Active threshold and was on a steeply falling trend. Observed fluence at 17/0000 UTC of 7.28e6 integrated pfu.

Four-Day Space Weather Forecast Summary

Solar Activity: Low to Moderate activity is now expected, with a slight chance of isolated X-class flares. These mainly from the new region close to the southeast limb which produced an

Issued by Met Office Space Weather Advisor, Tel: +44 (0) 330 135 4254 Email: moswoc@metoffice.gov.uk

© Crown Copyright 2024. All Rights Reserved.

X2.9 flare on 15 May. The rest of the regions currently on the visible disc are smaller and simpler, with isolated M-flares possible from these. A new region may rotate onto the northeast disc later Day 1 or Day 2 (17-18 May) with no other significant regions expected to return or rotate onto the disc during the period.

Solar Wind / Geomagnetic Activity: A glancing CME arrival is possible later Day 1 (17 May) from a filament eruption on the northeast disc on 14 May. Solar winds are expected to remain at background levels, but close to slightly elevated levels. Chance of some further enhancement from uncertain CME arrival. Activity is expected to be mainly Quiet (Kp 0-2) with Unsettled (Kp 3) intervals. Chance of an Active interval and a slight chance of a G1/Minor Geomagnetic storm intervals from any CME glances, mainly later Day 1 (17 May).

Energetic Particles / Solar Radiation: GOES-18 proton flux is expected to continue slowly easing, likely remaining below the S1/Minor radiation storm level. There is the potential for enhancements from any further significant flares from Ex-AR3664 on Day 1 (17 May), as the protons can still be directed towards Earth along the Parker Spiral. High energy electron flux reduced to background following CME arrival but forecast to rise once again into the period as CME influences eases, bringing a chance of Active fluence Days 2-4 (18-20 May), however this is low confidence due to uncertain CME arrival later Day 1.

Issued by Met Office Space Weather Advisor, Tel: +44 (0) 330 135 4254 Email: moswoc@metoffice.gov.uk

© Crown Copyright 2024. All Rights Reserved.

Figure 1. Solar Analysis Valid 17/0700 UTC.



Key: Filament ___ , Prominence ___ , Magnetic Field Line - - - , Polarity +/-, Coronal Holes: Grey shaded area CHxx +/-, Sunspot groups 25xx - Mt Wilson α-β-βγ-βγδ and Zurich-McIntosh Axx etc.

Issued by Met Office Space Weather Advisor, Tel: +44 (0) 330 135 4254 Email: moswoc@metoffice.gov.uk

© Crown Copyright 2024. All Rights Reserved.

Geomagnetic Storms:

The CMEs from Ex-AR3664 have now all passed Earth, with any further CMEs from this departed region expected to miss the Earth. There is the potential for a CME to partially arrive at Earth on Day 1 (17 May). This feature is associated with a filament eruption observed leaving the northeast disc on 14 May. While anticipated to pass largely north and behind Earth, glancing interactions are possible later Day 1, with associated enhancement potentially persisting overnight into Day 2 (18 May).

Solar winds have returned to background levels. In the absence of further CME enhancements, winds are expected to remain at background levels, but close to slightly elevated levels (350-400 km/s) with no notable coronal holes expected. Solar winds may see moderate enhancement to 450-500 km/s should any CME arrivals occur, this most likely Day 1 into Day 2 (17-18 May). Activity is expected to be mainly Quiet (Kp 0-2) with Unsettled (Kp 3) intervals. Chance of an Active interval and a slight chance of a G1/Minor Geomagnetic storm intervals from any CME glances, mainly later Day 1.

Geo-Magnetic Storm Probability (Exceedance)	Level	Past 24 Hours (Yes/No)	Day 1 (00-24 UTC) (%)	Day 2 (00-24 UTC) (%)	Day 3 (00-24 UTC) (%)	Day 4 (00-24 UTC) (%)
Minor or Moderate	G1 to G2	No	10	5	1	1
Strong	G3	No	1	1	1	1
Severe	G4	No	1	1	1	1
Extreme	G5	No	1	1	1	1

Geomagnetic Activity - Earthbound Coronal Mass Ejections

Date/time 21.5R (UTC)	Halo: Full or Partial	Source	Source Location	Estimated Speed	Estimated Arrival Time	Comments
14/1526	Partial	Filament eruption	NE	685 km/s	17/1800 UTC +/- 6hrs	Low confidence. Glancing blow.

Issued by Met Office Space Weather Advisor, Tel: +44 (0) 330 135 4254 Email: moswoc@metoffice.gov.uk

© Crown Copyright 2024. All Rights Reserved.

Figure 2: MOSWOC Ensemble showing possible glancing CME enhancement on 17 May.



Radio Blackouts - X-Ray Flares:

Low to Moderate activity is expected, with a slight chance of isolated High activity. The large and complex region that continues to rotate onto the southeast disc is considered to be the most likely source for X-class flares, having produced an X2.9 flare on 15 May. This region is currently difficult to analyse due to the restricted viewing angle, with a fuller understanding of the flare risk likely later in the coming 24 hours as this region continues rotating onto the visible disc. Recent flare activity has been mostly from AR3679, which is approaching the south central disc, with a peak C8.7 flare observed at 17/0146 UTC.

There are currently 14 sunspots on the visible disc, 13 of which are numbered. Any further activity from Ex-AR3664 is likely to be completely obscured as it has rotated onto the far-side. The largest region on the disc is AR3685 (Ehi/Beta), which lies close to the southeast limb. This region is composed of a number of large spots which may be a closely connected collection of different regions, magnetic looping is visible across the region and as such it remains classed as one large sunspot group. A new region continues to emerge from the southeast limb, just to the north of AR3685. At present it looks to be a single moderately large spot and is initially classed as an Hax/Alpha type. AR3679 (Esi/Beta-Gamma) is the second largest region, currently approaching the south central disc. It has also has been the source of most of the recent flare activity, with some further trailer spot emergence evident. AR3683 (Dsi/Beta-Gamma) in the southwest is of modest magnetic complexity and continues to show spot development/emergence. The other sunspot regions are generally smaller, simpler and stable or in slight decay.

Other than the emerging unnumbered region in the vicinity of AR3685, a further region may rotate onto the disc later Day 1 or Day 2 (17-18 May). Other significant regions may approach the eastern disc by the end of the period. As such, the flare risks have been maintained.

Issued by Met Office Space Weather Advisor, Tel: +44 (0) 330 135 4254 Email: moswoc@metoffice.gov.uk

© Crown Copyright 2024. All Rights Reserved.

X Ray Flares	Level	Past 24 Hours (Yes/No)	Day 1 (00-24 UTC)	Day 2 (00-24 UTC)	Day 3 (00-24 UTC)	Day 4 (00-24 UTC)
Probability			(%)	(%)	(%)	(%)
Active	R1-R2 M Class	No	60	60	60	60
Very Active	R3 to R5 X Class	No	10	10	10	10

Solar Radiation Storms - (High Energy Protons):

Recent S1/Minor Radiation Storm conditions were the result of the M6.6 flare of Ex-AR3664 on the southwest limb on 13 May and further contributions from the flaring of this region through 14-15 May, with at least four X-flares occurring.

Proton flux continues to decline and is expected to remain below S1/Minor storm levels, although there is a chance that should any large flares occur from Ex-AR3664 on Day 1 (17 May) that protons could be directed towards Earth through the Parker Spiral and produce some Minor radiation storming. While there are a number of small sunspot regions of the Earth-facing disc none of these are currently considered candidates for proton rich flares, with the potential for proton storms decreasing notably by Day 2 (18 May) as Ex-AR3664 moves further onto the far-side.

Radiation Storms	Level (cm^{-2} sr^{-1} s^{-1})	Past 24 Hours (Yes/No)	Day 1 (00-24 UTC)	Day 2 (00-24 UTC)	Day 3 (00-24 UTC)	Day 4 (00-24 UTC)
Probability (Exceedance)			(%)	(%)	(%)	(%)
Active	≥ S1	Yes	30	20	10	10
Very Active	≥ S3 *	No	1	1	1	1

* S3 ≥ 10 MeV ≥ 1000 pfu and / or ≥ 50 MeV ≥ 10 pfu. (pfu = $cm^{-2}sr^{-1}s^{-1}$)

Issued by Met Office Space Weather Advisor, Tel: +44 (0) 330 135 4254 Email: moswoc@metoffice.gov.uk

© Crown Copyright 2024. All Rights Reserved.

High Energy Electrons Event (≥ 2MeV):
High energy electron flux, as observed by GOES-16 has been at largely background levels following the CME arrival early on 16 May. This trend is most likely to persist in the short term, however there is the potential for electron flux to rise and potentially reach persistently High levels, with lower orbits having been highly charged by CME arrivals over previous days. A continued rise in flux levels from later Day 1 (17 May) onwards is expected although confidence is reduced due to the potential for a glancing CME arrival later on Day 1 (17 May), which may again reduce the observed flux.

The associated 24 hour fluence is expected to see a general falling trend near term, before likely retuning to a rising trend later Day 1 (17 May) onwards, perhaps approaching or crossing the Active threshold Days 2-4 (18-20 May). MOSWOC REFM providing reasonable guidance Day 1 although perhaps under representing the risk of reaching the Active threshold later in the period.

GEO Electron Environment	Level (cm^{-2} sr^{-1} day^{-1})	Past 24 Hours (Yes/No)	Day 1 (00-24 UTC)	Day 2 (00-24 UTC)	Day 3 (00-24 UTC)	Day 4 (00-24 UTC)
Probability (Exceedance)			(%)	(%)	(%)	(%)
Active	≥ 2 MeV ≥ $1x10^8$	No	10	20	30	30
Very Active	≥ 2 MeV ≥ $1x10^9$	No	1	1	1	1

Figure 3: MOSWOC REFM output.



Issued by Met Office Space Weather Advisor, Tel: +44 (0) 330 135 4254 Email: moswoc@metoffice.gov.uk

© Crown Copyright 2024. All Rights Reserved.